

The Lie derivative and Noether's theorem on the aromatic bicomplex

Adrien Laurent¹

July 16, 2023

Abstract

The aromatic bicomplex is an algebraic tool based on aromatic Butcher-trees and used in particular for the explicit description of volume-preserving affine-equivariant numerical integrators. To further understand the links between geometric numerical integration and variational calculus, as well as to describe more precisely the form of volume-preserving methods, we define and study the Lie derivative and the concept of symmetries on the aromatic bicomplex. We give an aromatic formulation of Noether's theorem and adapt the approach in the context of numerical volume-preservation.

Keywords: aromatic bicomplex, aromatic forms, Euler-Lagrange complex, Noether's theorem, aromatic Lie derivative, solenoidal forms, volume-preservation, geometric numerical integration.

AMS subject classification (2020): 58E30, 58J10, 05C05, 41A58, 37M15, 58A12.

1 Introduction

The search for an affine-equivariant volume-preserving method is one of the central open problems of geometric numerical integration. Such an integrator takes the form of an aromatic Butcher-series method [37]. While B-series [10, 21] (see also the textbooks [20, 11, 12] and the review [36]) describe the Taylor expansion of the flow of ordinary differential equations and of a large class of their numerical approximations, aromatic B-series were introduced in [24, 16] specifically for the study of volume preservation (see also [26, 35, 37, 6, 18, 7, 33]) as B-series methods cannot preserve volume in general. We mention that finding a volume-preserving aromatic B-series method is the first step toward an exotic aromatic S-series method [30, 31, 27, 8, 29] that exactly preserves the invariant measure of ergodic stochastic differential equations as the algebraic conditions are similar (see, for instance, [1]).

The recent work [28] introduces new tools from the calculus of variations, such as the aromatic bicomplex (see also [2, 3, 40] and references therein), and these tools yield valuable insight of the form of the Taylor expansion of a volume-preserving methods. In particular, it shows that aromatic Runge-Kutta methods do not preserve volume in general, while aromatic exponential methods are promising starting points. To further understand the form of a volume-preserving method, the present work defines new tools on the aromatic bicomplex such as the Lie derivative, different concepts of symmetries, and Noether's theory in the aromatic context.

Let τ, γ be linear combinations of trees, the Lie derivative of γ in the direction of τ is

$$\mathcal{L}_\tau \gamma = \left. \frac{d}{d\varepsilon} \right|_{\varepsilon=0} [(\bullet + \varepsilon\tau) \triangleright \gamma],$$

¹Department of Mathematics, University of Bergen, Bergen, Norway. Adrien.Laurent@uib.no.

where \triangleright is the substitution law of B-series [20, 15, 13] (see also [6, 9] for the substitution of aromatic and exotic aromatic series). The Lie derivative is also the pre-Lie version of the substitution law on B-series. It appears under the name pre-Lie insertion product in [43, 34, 44] for the study of the freeness of the pre-Lie insertion algebra. In the calculus of variations, the Lie-derivative defines symmetries, that are perturbations that leave the input unchanged at first order. This leads to the Noether theorem, that draws links between symmetries and conservation laws.

This paper gives a general definition of the Lie derivative on aromatic forms by using the framework given by the aromatic bicomplex. This allows us to define symmetries, to write an aromatic version of the Noether theorem, and to draw further links between variational calculus and numerical volume-preservation. We give a concise introduction in Section 2 of the aromatic bicomplex and its properties, while Section 3 is devoted to the general definition of the Lie derivative, of the different symmetries, and the statement of the aromatic Noether theorem. We then adapt the new approach and results to the study of volume preserving methods.

2 Preliminaries on the aromatic bicomplex

In this section, we give a concise definition of the necessary tools and concepts required for the definition of the aromatic Euler-Lagrange complex. The notations and vocabulary are chosen to match with the literature of variational calculus, as we shall draw bridges between numerical analysis and variational calculus in Section 3. We skip the technical details on the Euler and homotopy operators and refer the reader to [28] for more details.

2.1 Aromatic forms and their derivatives

While aromatic trees represent vector fields, we use aromatic forests to represent specific classes of homogeneous tensors and forms. This allows in particular to translate the technicalities of the infinite jet bundle $J^\infty(\mathbb{R}^d)$ into straightforward combinatorics.

Definition 2.1. *Let V be a finite set of nodes, split into vertices V^\bullet and covertices V° , and $E \subset V \times V$ a set of oriented edges. The covertices are numbered from 1 to p , while the vertices are indistinguishable. Each node in V is the source of exactly one edge, except the roots that have no outgoing edges, that we order and number from 1 to n . Any connected component of such a graph either has exactly one root, and is called a tree, or does not have a root, and is called an aroma. We call aromatic forests such graphs, up to equivalence of graphs that preserve the numbering of the covertices and the roots. We write $\mathcal{F}_{n,p}$ the set of aromatic forests with n roots and p covertices and $\mathcal{F}_n = \mathcal{F}_{n,0}$. The number of nodes $|\gamma|$ of an aromatic forest γ is called the order of γ .*

In the spirit of differential geometry, we alternatize aromatic forests to obtain aromatic forms using the wedge projection operator.

Definition 2.2. *For $\gamma \in \mathcal{F}_{n,p}$, let \mathcal{S}_n^\bullet (resp. \mathcal{S}_p°) be the set of permutations of the roots of γ (resp. the covertices of γ). The roots and covertices wedges of γ are*

$$\wedge^\bullet \gamma = \frac{1}{n!} \sum_{\sigma \in \mathcal{S}_n^\bullet} \varepsilon(\sigma) \sigma \gamma, \quad \wedge^\circ \gamma = \frac{1}{p!} \sum_{\sigma \in \mathcal{S}_p^\circ} \varepsilon(\sigma) \sigma \gamma,$$

where $\varepsilon(\sigma)$ is the signature of the permutation σ . The wedge operator is $\wedge = \wedge^\bullet \wedge^\circ = \wedge^\circ \wedge^\bullet$. We extend the wedge operations to $\text{Span}(\mathcal{F}_{n,p})$ by linearity and we denote the set of aromatic forms $\Omega_{n,p} = \wedge \text{Span}(\mathcal{F}_{n,p})$ and $\Omega_n = \Omega_{n,0}$.

Example. Let $\gamma_1 = \bullet \dot{\downarrow} \in \mathcal{F}_2$, $\gamma_2 = \begin{array}{c} \textcircled{2} \\ \textcircled{1} \end{array} \in \mathcal{F}_{0,2}$, $\gamma_3 = \textcircled{1} \textcircled{2} \in \mathcal{F}_{2,2}$, then

$$\wedge \gamma_1 = \frac{1}{2}(\bullet \dot{\downarrow} - \dot{\downarrow} \bullet) \in \Omega_2, \quad \wedge \gamma_2 = 0, \quad \wedge \gamma_3 = \frac{1}{2}(\textcircled{1} \textcircled{2} - \textcircled{2} \textcircled{1}).$$

The operations on the variational bicomplex make use of differentiations in the infinite jet bundle. In the aromatic context, we replace the differentiations by the operations of grafting and replacing nodes. The horizontal derivative is defined using grafting operations, while the vertical derivative uses the replacing operation. The sign change in the definition of the total derivative is explained in [28, Rk. 2.8]. We mention that the horizontal and vertical derivatives were used in a different context on $\Omega_1 = \text{Span}(\mathcal{F}_1)$ respectively in [16, 24] for d_H and in [18] for d_V .

Definition 2.3. Let $\gamma \in \mathcal{F}_{n,p}$, r a root of γ , and $u \in V$ (possibly equal to r), then $D^{r \rightarrow u} \gamma$ returns a copy of γ where the node r is now a predecessor of u . The operator $D^r \gamma = \sum_{u \in V} D^{r \rightarrow u} \gamma$ grafts r to all possible nodes. Let $\gamma \in \mathcal{F}_{n,p}$ and $v \in V^\bullet$, then $\gamma_{v \rightarrow \textcircled{k}}$ is the forest obtained by replacing the node v by a new covertex \textcircled{k} . Similarly, $\gamma_{\textcircled{k} \rightarrow \tau}$ is the linear combination of forests obtained by replacing the covertex \textcircled{k} by the tree τ and grafting the predecessors of \textcircled{k} to the nodes of τ in all possible ways. The horizontal, vertical, and total derivatives of $\gamma \in \mathcal{F}_{n,p}$ are

$$d_H \gamma = D^{r_n} \gamma, \quad d_V \gamma = \wedge \sum_{v \in V^\bullet} \gamma_{v \rightarrow \textcircled{p+1}}, \quad d\gamma = (-1)^{n+p} d_H \gamma + d_V \gamma.$$

We extend d_H and d_V on $\Omega_{n,p}$ by linearity into $d_H: \Omega_{n,p} \rightarrow \Omega_{n-1,p}$ and $d_V: \Omega_{n,p} \rightarrow \Omega_{n,p+1}$, with the convention $d_H \gamma = 0$ if $\gamma \in \Omega_{0,p}$.

Example. Consider $\gamma_1 = \bullet \in \Omega_1$, $\gamma_2 = \wedge \bullet \dot{\downarrow} \in \Omega_2$, and $\gamma_3 = \wedge \bullet \textcircled{1} \in \Omega_{2,1}$, then

$$d_H \gamma_1 = \circlearrowleft, \quad d_H \gamma_2 = \frac{1}{2}(\circlearrowleft \bullet + \bullet \circlearrowleft - \circlearrowleft \dot{\downarrow} - \dot{\downarrow} \circlearrowleft), \quad d_H \gamma_3 = \frac{1}{2}(\textcircled{1} \bullet + \bullet \textcircled{1} - \circlearrowleft \textcircled{1} - \textcircled{1} \circlearrowleft),$$

$$d_V \gamma_1 = \textcircled{1}, \quad d_V \gamma_2 = \wedge \textcircled{1} \dot{\downarrow} + \wedge \bullet \textcircled{1} + \wedge \bullet \textcircled{1}, \quad d_V \gamma_3 = \wedge \textcircled{2} \textcircled{1} = \frac{1}{2}(\textcircled{2} \textcircled{1} - \textcircled{1} \textcircled{2}).$$

The derivatives on aromatic forms naturally form complexes as justified by the following result. Note that the horizontal and vertical derivatives commute, while their equivalents in variational calculus anticommute [3].

Proposition 2.4 ([28]). *The derivatives satisfy*

$$d_H^2 = 0, \quad d_V^2 = 0, \quad d_V d_H = d_H d_V, \quad d^2 = 0.$$

2.2 The aromatic bicomplex and the Euler-Lagrange complex

The object of ultimate interest in variational calculus is the Euler-Lagrange complex, which requires defining the augmented bicomplex first. The interior Euler operator I is given for $\gamma \in \Omega_{0,1}$ by the combination of forms obtained by unplugging all the predecessors of the covertex of γ , plugging back the predecessors on all the vertices in all possible ways, and multiplying by -1 if the number of predecessors is odd. For instance, we have

$$I \textcircled{1} = 0, \quad I \textcircled{\circlearrowleft} = \textcircled{\circlearrowleft}, \quad I \textcircled{\bullet \dot{\downarrow}} = \textcircled{\bullet \dot{\downarrow}}, \quad I \textcircled{\bullet \textcircled{1}} = -\textcircled{\bullet \textcircled{1}}, \quad I \textcircled{\circlearrowleft \textcircled{1}} = -\textcircled{\circlearrowleft \textcircled{1}}.$$

For the sake of simplicity, we refer to [28, Sect. 4.1] for the precise definition of $I: \Omega_{0,p} \rightarrow \Omega_{0,p}$. The variational derivative $\delta_V = I \circ d_V$ and the interior Euler operator I satisfy

$$I^2 = I, \quad Id_H = 0, \quad \delta_V^2 = 0.$$

In particular, I is a projection on $\mathcal{I}_p = I(\Omega_{0,p})$, the aromatic equivalent of the space of source forms.

The augmented aromatic bicomplex is the diagram drawn in Figure 1 that displays the interactions between the different spaces of aromatic forms. The aromatic bicomplex can be seen as a generalised subcomplex of the variational bicomplex [2, 3] that focuses on specific classes of homogeneous forms and is independent of the dimension of the problem (see [28, Rk. 2.8]).

$$\begin{array}{ccccccc}
& & \vdots & & \vdots & & \vdots & & \vdots & & \\
& & d_V \uparrow & & d_V \uparrow & & d_V \uparrow & & \delta_V \uparrow & & \\
\dots & \xrightarrow{d_H} & \Omega_{2,2} & \xrightarrow{d_H} & \Omega_{1,2} & \xrightarrow{d_H} & \Omega_{0,2} & \xrightarrow{I} & \mathcal{I}_2 & \longrightarrow & 0 \\
& & d_V \uparrow & & d_V \uparrow & & d_V \uparrow & & \delta_V \uparrow & & \\
\dots & \xrightarrow{d_H} & \Omega_{2,1} & \xrightarrow{d_H} & \Omega_{1,1} & \xrightarrow{d_H} & \Omega_{0,1} & \xrightarrow{I} & \mathcal{I}_1 & \longrightarrow & 0 \\
& & d_V \uparrow & & d_V \uparrow & & d_V \uparrow & & \delta_V \nearrow & & \\
\dots & \xrightarrow{d_H} & \Omega_2 & \xrightarrow{d_H} & \Omega_1 & \xrightarrow{d_H} & \Omega_0 & & & & \\
& & \uparrow & & \uparrow & & \uparrow & & & & \\
& & 0 & & 0 & & 0 & & & &
\end{array}$$

Figure 1: The augmented aromatic bicomplex.

The bottom row of the aromatic bicomplex is similar to the De Rham complex. It extends into the edge complex (2.1), called the aromatic Euler-Lagrange complex.

$$\dots \xrightarrow{d_H} \Omega_2 \xrightarrow{d_H} \Omega_1 \xrightarrow{d_H} \Omega_0 \xrightarrow{\delta_V} \mathcal{I}_1 \xrightarrow{\delta_V} \mathcal{I}_2 \xrightarrow{\delta_V} \dots \quad (2.1)$$

We introduce some vocabulary to further motivate the importance of the aromatic Euler-Lagrange complex. While we call aromatic forms the elements of $\Omega_{n,p}$, the space Ω_1 spanned by aromatic trees represent both vector fields and differential forms in variational calculus, so that we shall call the elements of Ω_1 aromatic vector fields or aromatic forms depending on the context. The elements of Ω_0 spanned by multi-aromas represent Lagrangians in variational calculus and volume forms in the context of volume-preservation, so that we call its elements aromatic Lagrangians. The source forms in \mathcal{I}_1 represent differential equations. The Euler-Lagrange complex (2.1) is the rigorous implementation of the following diagram.

$$\dots \xrightarrow{\text{curl}} \text{vector fields} \xrightarrow{\text{divergence}} \text{Lagrangians} \xrightarrow{\text{Euler-Lagrange}} \text{diff. equations} \xrightarrow{\text{Helmotz}} \dots$$

The crucial property of the aromatic bicomplex is its exactness. We refer the reader to [28] for the proof of the exactness and the explicit expressions of the associated homotopy operators.

Theorem 2.5 ([28]). *The horizontal and vertical sequences of the augmented aromatic bicomplex and the aromatic Euler-Lagrange complex (2.1) are exact, that is, there exist homotopy operators*

$$h_H: \Omega_{n,p} \rightarrow \Omega_{n+1,p}, \quad h_V: \Omega_{n,p} \rightarrow \Omega_{n,p-1}, \quad \mathfrak{h}_H: \Omega_{0,p} \rightarrow \Omega_{1,p}, \quad \mathfrak{h}_V: \mathcal{I}_p \rightarrow \mathcal{I}_{p-1},$$

such that the following identities hold:

$$\gamma = (d_H h_H + h_H d_H)\gamma, \quad \gamma \in \Omega_{n,p}, \quad n \geq 1, \quad p \geq 0, \quad (2.2)$$

$$\gamma = (d_V h_V + h_V d_V)\gamma, \quad \gamma \in \Omega_{n,p}, \quad n \geq 0, \quad p \geq 1, \quad (2.3)$$

$$\gamma = (d_H h_H + h_V \delta_V)\gamma, \quad \gamma \in \Omega_0,$$

$$\gamma = (I + d_H \mathfrak{h}_H)\gamma, \quad \gamma \in \Omega_{0,p}, \quad p \geq 1, \quad (2.4)$$

$$\gamma = (\delta_V \mathfrak{h}_V + \mathfrak{h}_V \delta_V)\gamma, \quad \gamma \in \mathcal{I}_p, \quad p \geq 1.$$

In particular, if one is interested in the description of all aromatic vector fields $\gamma \in \Omega_1$ of vanishing divergence $d_H \gamma = 0$, then Theorem 2.5 states that $\gamma = d_H \eta \in \text{Im}(d_H)$. The problem of finding a volume-preserving numerical method translates via backward error analysis [20] into the precise description of $\text{Ker}(d_H|_{\Omega_1})$, so that the Euler-Lagrange complex becomes a strong tool in this context. We further describe $\text{Ker}(d_H|_{\Omega_1})$ using symmetries and Noether's theorem.

3 Noether's theory on the aromatic bicomplex

In this section, we define the Lie derivative of aromatic forms, study its properties, and use it to rewrite the Noether theorem in the context of the aromatic bicomplex. We then apply these new tools in the context of volume-preservation.

3.1 The aromatic Lie derivative

In the spirit of the differential geometry literature [32], we use a Cartan formula to define the aromatic Lie derivative on aromatic forms. We then present the different properties of the Lie derivative.

Definition 3.1. *For $\gamma \in \mathcal{F}_{n,p}$, the contraction of γ in direction of $\tau \in \mathcal{F}_1$ is*

$$i_\tau \gamma = p \gamma_{\textcircled{p} \rightarrow \tau},$$

where $i_\tau \gamma = 0$ if $p = 0$. We extend the contraction in $i_\tau: \Omega_{n,p} \rightarrow \Omega_{n,p-1}$ for $\tau \in \Omega_1$ by linearity. For an aromatic form $\gamma \in \Omega_{n,p}$ and an aromatic vector field $\tau \in \Omega_1$, the aromatic Lie derivative of γ in the direction of τ is given by the Cartan formula:

$$\mathcal{L}_\tau \gamma = (d i_\tau + i_\tau d)\gamma. \quad (3.1)$$

Thanks to the homotopy identities of Theorem 2.5, the Lie derivative satisfies the following identities.

Proposition 3.2. *For $\tau \in \Omega_1$, the Lie derivative $\mathcal{L}_\tau \gamma: \Omega_{n,p} \rightarrow \Omega_{n,p}$ satisfies*

$$\mathcal{L}_\tau \gamma = (d_V i_\tau + i_\tau d_V)\gamma, \quad \gamma \in \Omega_{n,p}, \quad (3.2)$$

$$\mathcal{L}_\tau \gamma = (d_H H_\tau + H_\tau d_H)\gamma, \quad \gamma \in \Omega_n, \quad n > 0,$$

$$\mathcal{L}_\tau \gamma = (d_H \mathfrak{H}_\tau + i_\tau \delta_V)\gamma, \quad \gamma \in \Omega_0,$$

where $H_\tau = \mathcal{L}_\tau h_H$ and $\mathfrak{H}_\tau = \mathcal{L}_\tau \mathfrak{h}_H$ are the Lie homotopy operators.

Proof. As d_H and i_τ commute, replacing $d = (-1)^{n+p}d_H + d_V$ in (3.1) gives (3.2). We then deduce from the expression (3.2) that $\mathcal{L}_\tau\gamma: \Omega_{n,p} \rightarrow \Omega_{n,p}$. For $n > 0$ and $p = 0$, the horizontal homotopy identity (2.2) yields

$$\mathcal{L}_\tau\gamma = i_\tau d_V\gamma = i_\tau d_V d_H h_H\gamma + i_\tau d_V h_H d_H\gamma = d_H H_\tau\gamma + H_\tau d_H\gamma,$$

where we used that d_H commutes with i_τ and d_V and that d_V commutes with h_H . Similarly, the augmented horizontal homotopy identity (2.4) yields the expression of \mathcal{L}_τ on Ω_0 . \square

Remark 3.3. *The vertical homotopy operator is linked to the contraction by $i_\bullet\gamma = |\gamma| h_V\gamma$. Thus, equations (2.3) and (3.2) yield*

$$\mathcal{L}_\bullet\gamma = |\gamma|\gamma.$$

The equivalent of this property is used in [38] with planar forests to obtain the expansion of the Grossman-Larsson exponential.

On the geometrical side, the Lie derivative naturally realises a Lie algebra structure.

Proposition 3.4. *For $\tau_1, \tau_2 \in \Omega_1$, define the commutator on aromatic vector fields by*

$$[[\tau_1, \tau_2]] := \mathcal{L}_{\tau_1}\tau_2 - \mathcal{L}_{\tau_2}\tau_1.$$

Then, $(\Omega_1, [[\cdot, \cdot]])$ is a Lie algebra and \mathcal{L} is a Lie algebra representation, that is, for $\gamma \in \Omega_n$,

$$[\mathcal{L}_{\tau_1}, \mathcal{L}_{\tau_2}]\gamma := \mathcal{L}_{\tau_1}\mathcal{L}_{\tau_2}\gamma - \mathcal{L}_{\tau_2}\mathcal{L}_{\tau_1}\gamma = \mathcal{L}_{[[\tau_1, \tau_2]]}\gamma.$$

Proof. The Jacobi identity for the bracket $[[\cdot, \cdot]]$ is a consequence of the pre-Lie property of the Lie derivative [43, 34, 44], extended straightforwardly to aromatic trees. Define for $v \in V$ a node of γ , $\mathcal{L}_\tau^v\gamma = i_\tau(\gamma_{v \rightarrow \textcircled{1}})$. Then, we observe

$$\mathcal{L}_{\tau_1}\mathcal{L}_{\tau_2}\gamma = \sum_{\substack{v, w \in V \\ v \neq w}} \mathcal{L}_{\tau_1}^w \mathcal{L}_{\tau_2}^v \gamma + \mathcal{L}_{\mathcal{L}_{\tau_1}\tau_2}\gamma.$$

As $\mathcal{L}_{\tau_1}^w$ and $\mathcal{L}_{\tau_2}^v$ commute, a similar expression for $\mathcal{L}_{\tau_2}\mathcal{L}_{\tau_1}\gamma$ yields the result. \square

Thanks to equation (3.2), the classical geometric properties of the Lie derivative extend to aromatic forms.

Proposition 3.5. *For $\tau \in \Omega_1$, the Lie derivative \mathcal{L}_τ commutes with the derivatives d_H, d_V . In particular, we have $[\mathcal{L}_\tau, d] = 0$. Moreover, for $\mu \in \Omega_0, \tau \in \Omega_1$, and $\gamma \in \Omega_n$, the following product rule holds,*

$$\mathcal{L}_\tau(\mu\gamma) = \mu(\mathcal{L}_\tau\gamma) + (\mathcal{L}_\tau\mu)\gamma.$$

Proof. The Lie derivative commutes with d_H as d_H commutes with d_V and i_τ . Using Proposition (2.4) yields

$$\mathcal{L}_\tau d_V = d_V i_\tau d_V = d_V \mathcal{L}_\tau.$$

The product rule is a consequence of the product rule for d_V , that is,

$$d_V(\mu\gamma) = \mu(d_V\gamma) + (d_V\mu)\gamma.$$

Hence the result. \square

The two Hopf algebra structures on standard B-series are associated to the composition and substitution laws [17, 15, 13] (see also [6, 8, 42]). The associated pre-Lie laws on Ω_1 are the grafting product, given for $\tau, \gamma \in \Omega_1$, r_τ the root of τ , and V_γ the vertices of γ by

$$\tau \curvearrowright \gamma = \sum_{v \in V_\gamma} D^{r_\tau \rightarrow v}(\tau\gamma),$$

and the insertion product [43, 34, 44], that coincides with the Lie derivative $\mathcal{L}_\tau \gamma$. We mention that the dual of the grafting product is also called Lie derivative (though it differs from $\mathcal{L}_\tau \gamma$) in [19] (see also [20, Sec. IX.9.1]) and is used for computing the modified equation of a B-series method in terms of trees. The two pre-Lie structures interact according to the following identity, which is the pre-Lie version of the compatibility relation between the laws of composition and substitution of aromatic B-series. For $\tau_1, \tau_2 \in \Omega_1$ and $\gamma \in \Omega_n$, we have

$$\mathcal{L}_{\tau_1}(\tau_2 \curvearrowright \gamma) = (\mathcal{L}_{\tau_1} \tau_2) \curvearrowright \gamma + \tau_2 \curvearrowright (\mathcal{L}_{\tau_1} \gamma). \quad (3.3)$$

Remark 3.6. Thanks to [18], it is known that Ω_1 is freely generated from $\{\bullet\}$ by the operations \curvearrowright and d_H . Thus, we deduce that $\Omega_{n,p}$ is freely generated from $\{\bullet, \textcircled{\bullet}, \textcircled{\circ}, \dots\}$ by the operations \curvearrowright , d_H , and concatenation. For example, the aromatic Lagrangian $\mathfrak{Q} = \textcircled{\circ}$ decomposes into

$$\mathfrak{Q} = d_H(\bullet \curvearrowright \bullet) - \bullet \curvearrowright d_H \bullet.$$

This allows us to write an alternative expression of the Lie derivative by replacing in each term one node \bullet by τ in all possible ways:

$$\mathcal{L}_\tau \mathfrak{Q} = d_H(\tau \curvearrowright \bullet) + d_H(\bullet \curvearrowright \tau) - \tau \curvearrowright d_H \bullet - \bullet \curvearrowright d_H \tau.$$

The substitution law \triangleright of aromatic vector fields can be rewritten in terms of the Lie derivative in the spirit of [41]. We first extend \mathcal{L} on $\mathcal{U}(\Omega_1) \times \mathcal{U}(\Omega_1)$, where $\mathcal{U}(\Omega_1)$ is the universal enveloping algebra of Ω_1 equipped with the symmetric product \cdot and the shuffle coproduct Δ . Monomials in $\mathcal{U}(\Omega_1)$ are also called clumped forests in [9] (see also [6]), and in particular we have in $\mathcal{U}(\Omega_1)$:

$$(\textcircled{\circ} \textcircled{\bullet}) \cdot (\textcircled{\circ} \bullet) = \frac{1}{2}(\textcircled{\circ} \textcircled{\bullet})(\textcircled{\circ} \bullet) + \frac{1}{2}(\textcircled{\circ} \bullet)(\textcircled{\circ} \textcircled{\bullet}) \neq (\textcircled{\circ} \textcircled{\bullet} \textcircled{\bullet}) \dots$$

The Lie derivative is extended on $\mathcal{U}(\Omega_1)$ using the following rules from [41]:

$$\mathcal{L}_{\mathbb{1}} \gamma_1 = \gamma_1, \quad \mathcal{L}_{\tau \gamma_2} \gamma_1 = \mathcal{L}_\tau \mathcal{L}_{\gamma_2} \gamma_1 - \mathcal{L}_{\mathcal{L}_\tau \gamma_2} \gamma_1, \quad \mathcal{L}_{\gamma_3}(\gamma_1 \cdot \gamma_2) = (\mathcal{L}_{\gamma_3^{(1)}} \gamma_1) \cdot (\mathcal{L}_{\gamma_3^{(2)}} \gamma_2),$$

where $\tau \in \Omega_1$, $\gamma_1, \gamma_2, \gamma_3 \in \mathcal{U}(\Omega_1)$, and we use Sweedler's notation.

Proposition 3.7. Let $\tau, \gamma \in \Omega_1$ and $\varepsilon > 0$, then the perturbation of γ by $\varepsilon \tau$ is

$$(\bullet + \varepsilon \tau) \triangleright \gamma = \mathcal{L}_{\exp(\varepsilon \tau)} \gamma, \quad \exp(\varepsilon \tau) = \mathbb{1} + \varepsilon \tau + \frac{\varepsilon^2}{2} \tau \cdot \tau + \frac{\varepsilon^3}{3!} \tau \cdot \tau \cdot \tau + \dots$$

3.2 Symmetries and divergence symmetries

The Lie derivative allows us to define the concepts of symmetries and divergence symmetries.

Definition 3.8. The aromatic vector field $\tau \in \Omega_1$ is a symmetry for the aromatic form γ if $\mathcal{L}_\tau \gamma = 0$. If there exists an aromatic form η such that $\mathcal{L}_\tau \gamma = d_H \eta$, we say that τ is a divergence symmetry for γ .

The simplest symmetries are also called solenoidal [28]. They are the object of ultimate interest in numerical volume-preservation.

Definition 3.9. *An aromatic vector field $\tau \in \Omega_1$ is solenoidal if it is a symmetry for the aromatic Lagrangian $\mathring{\circ}$.*

Thanks to the exactness of the aromatic bicomplex (see Theorem 2.5), the set of solenoidal forms is $d_H(\Omega_2)$. In particular, the simplest examples of solenoidal forms are

$$\begin{aligned}
2d_H \wedge \cdot \dot{\circ} &= \mathring{\circ} \cdot + \mathring{\circ} \cdot - \mathring{\circ} \dot{\circ} - \mathring{\vee}, \\
2d_H \wedge \cdot \dot{\circ} \dot{\circ} &= \mathring{\circ} \cdot + \mathring{\circ} \cdot + \mathring{\Delta} \cdot - \mathring{\vee} \dot{\circ} - \mathring{\vee} \dot{\circ} - \mathring{\circ} \dot{\circ} \dot{\circ}, \\
2d_H \wedge \cdot \mathring{\vee} &= \mathring{\circ} \cdot + 2\mathring{\circ} \cdot + \mathring{\vee} \dot{\circ} - 2\mathring{\vee} \dot{\circ} - \mathring{\vee} \dot{\circ} - \mathring{\circ} \mathring{\vee}, \\
2d_H \wedge \mathring{\circ} \dot{\circ} &= \mathring{\circ} \cdot + \mathring{\circ} \mathring{\circ} \cdot + \mathring{\circ} \mathring{\circ} \cdot - \mathring{\circ} \dot{\circ} - \mathring{\circ} \dot{\circ} - \mathring{\circ} \mathring{\vee}.
\end{aligned}$$

As the complexity of the calculation increases rapidly with the order, there is no known example of symmetry that is not solenoidal at the present time.

Let us state the aromatic formulation of the first variational formula, a central tool in the proof of Noether's theorem.

Proposition 3.10 (First variational formula). *Let $\gamma \in \Omega_0$ (respectively $\gamma \in \mathcal{I}_p$), then there exists an aromatic form $\eta \in \Omega_{1,1}$ (respectively $\eta \in \Omega_{1,p+1}$) such that*

$$d_V \gamma = \delta_V \gamma + d_H \eta. \quad (3.4)$$

In particular, let $\tau \in \Omega_1$ and $\gamma \in \Omega_0$, then there exists $\eta \in \Omega_1$ such that

$$\mathcal{L}_\tau \gamma = i_\tau \delta_V \gamma + d_H \eta. \quad (3.5)$$

Proof. Let $\gamma \in \Omega_0$ or $\gamma \in \mathcal{I}_p$ and let $\omega = d_V \gamma$. As $\omega - I\omega \in \text{Ker}(I)$, by horizontal exactness of the aromatic bicomplex (Theorem 2.5), there exists η such that

$$d_V \gamma = \omega = I\omega + (\omega - I\omega) = I\omega + d_H \eta.$$

Applying the contraction i_τ to (3.4) yields (3.5) as i_τ and d_H commute. \square

A first application of the first variational formula, in particular of equation (3.5), is that any $\tau \in \Omega_1$ is a divergence symmetry for $\gamma \in \text{Ker}(\delta_V|_{\Omega_0}) = d_H(\Omega_1)$. In particular, solenoidal forms (and more generally symmetries) are divergence symmetries. The elements with up to three nodes in $d_H(\Omega_1)$ are spanned by

$$\{ \mathring{\circ}, \mathring{\circ} + \mathring{\circ}, \mathring{\circ} + \mathring{\circ} \mathring{\circ}, \mathring{\circ} + \mathring{\circ} + \mathring{\Delta}, \mathring{\vee} + 2\mathring{\circ}, \mathring{\circ} + \mathring{\circ} \mathring{\circ} + \mathring{\circ} \mathring{\circ}, \mathring{\vee} - \mathring{\circ} \mathring{\circ}, \mathring{\circ} \mathring{\circ} \mathring{\circ} + 2\mathring{\circ} \mathring{\circ} \}$$

An alternative way to produce symmetries is the following consequence of Propositions 3.4 and 3.5.

Proposition 3.11. *If $\tau_1 \in \Omega_1$ is a divergence symmetry for $\gamma \in \Omega_0$, then for any $\tau_2 \in \Omega_1$, $[[\tau_1, \tau_2]]$ is a divergence symmetry for γ . If τ_1 is a symmetry for γ , then τ_1 is a symmetry for $\mathcal{L}_{\tau_2} \gamma$ if and only if $[[\tau_1, \tau_2]]$ is a symmetry for γ .*

Remark 3.12. *As the number of aromatic trees grows fast with the number of nodes, one is often interested in finding symmetries for specific differential systems. Let $f: \mathbb{R}^d \rightarrow \mathbb{R}^d$ be a smooth vector field, and F be the elementary differential map (see [28]). We call f -symmetry for γ an aromatic vector field $\tau \in \Omega_1$ such that $F(\mathcal{L}_\tau \gamma)(f) = 0$, and τ is f -solenoidal in the specific case where $\gamma = \circlearrowleft$. In particular, if we assume that $\operatorname{div}(f) = 0$, then $\tau = \circlearrowleft \cdot - \searrow \swarrow$ is a f -symmetry for \circlearrowleft . The assumption $\operatorname{div}(f) = 0$ does not create new f -solenoidal vector fields [28], but it does create new f -symmetries. In particular, a f -solenoidal aromatic vector field with $\operatorname{div}(f) = 0$ is a f -symmetry of any aroma $\gamma \in \Omega_0$ with a 1-loop. For example, $\tau = \circlearrowleft \cdot - \searrow \swarrow$ is a f -symmetry for \circlearrowleft , \circlearrowright , $\circlearrowleft \cdot$, $\circlearrowright \cdot$, $\searrow \swarrow$, $\swarrow \searrow$, \dots . The use of specific vector fields f gives rise to degeneracies, which makes it easier to find f -symmetries. We cite in particular the work [7, Sec. 4] that uses such degeneracies with f quadratic.*

3.3 The aromatic Noether theorem

The Noether theorem, published by Emmy Noether in 1918 in her paper *Invariante Variation-sprobleme* (see the english translation [39]), draws an explicit link between symmetries and conservation laws in the context of variational calculus.

Definition 3.13. *The aromatic vector field $\tau \in \Omega_1$ is a generator of a conservation law $\eta \in \Omega_1$ for the source form $\gamma \in \mathcal{I}_1$ if $i_\tau \gamma = d_H \eta$.*

We rewrite Noether's theorem in the context of aromatic forms.

Theorem 3.14 (Noether's theorem). *Let an aromatic Lagrangian $\gamma \in \Omega_0$ and an aromatic vector field $\tau \in \Omega_1$. Then, τ is a divergence symmetry of γ if and only if τ is the generator of a conservation law for $\delta_V \gamma$.*

Theorem 3.14 is a direct consequence of the first variational formula (3.5). In the context of variational calculus, a symmetry of the Lagrangian exactly corresponds to the preservation of a quantity, which typically leads to superfluous degrees of freedom (see, for instance, the textbooks [2, 4, 5]).

Example. *Let $\gamma \in d_H(\Omega_1)$, then as d_H and \mathcal{L}_τ commute, any aromatic vector field τ is a divergence symmetry for γ . The associated conservation law is $\eta = 0$.*

Remark 3.15. *An alternate formulation of Theorem 3.14 with source forms is the following. Define the natural Lie derivative on \mathcal{I}_p by $\mathcal{L}_\tau^\natural = I\mathcal{L}_\tau$. Assume that $\gamma \in \mathcal{I}_1$ satisfies $\delta_V \gamma = 0$ (i.e., γ is variational), then the aromatic vector field $\tau \in \Omega_1$ is a symmetry for γ if and only if $\delta_V i_\tau \gamma = 0$. This formulation of the Noether theorem is a consequence of the first variational formula (3.5) and the identities*

$$\mathcal{L}_\tau^\natural \gamma = (\delta_V i_\tau + I i_\tau \delta_V) \gamma, \quad \delta_V \mathcal{L}_\tau \gamma = \mathcal{L}_\tau^\natural d_V \gamma.$$

3.4 Application to numerical volume-preservation

The search for a volume-preserving integrator in the form of an aromatic B-series method, or equivalently of an affine-equivariant method [35, 37, 29], is an important open question of geometric numerical integration. It is known that there is no volume-preserving B-series method except the exact flow [16, 24], but the question for aromatic B-series methods is still open. We refer to the recent works [6, 7, 14, 28] for different approaches to the conjecture. In this

subsection, we apply the results on the Lie derivative, the symmetries, and the Noether theorem in the context of volume-preservation, and we rewrite the numerical open questions in a purely algebraic manner.

We denote $\overline{\Omega}_1$ the set of formal series of elements of Ω_1 graded by the number of nodes of the aromatic trees, also called aromatic B-series. The results presented in this paper extend straightforwardly to aromatic B-series. Thanks to Proposition 3.7 and [28, Thm.4.17], the problem of volume-preservation boils down to finding an aromatic B-series method (or class of methods) that has the following form:

$$(\bullet + \tau) \triangleright \gamma = \mathcal{L}_{\exp(\tau)}\gamma$$

where τ is solenoidal and $\gamma \in \overline{\Omega}_1$ is the B-series of the exact flow of $y' = f(y)$ (see [20, Chap. III]). In particular, the first terms of the expansion are

$$\mathcal{L}_{\exp(\tau)}\gamma = \gamma + \mathcal{L}_\tau\gamma + \frac{1}{2}(\mathcal{L}_\tau^2 - \mathcal{L}_{\mathcal{L}_\tau\tau})\gamma + \frac{1}{6}(\mathcal{L}_\tau^3 - 2\mathcal{L}_\tau\mathcal{L}_{\mathcal{L}_\tau\tau} - \mathcal{L}_{\mathcal{L}_\tau\tau}\mathcal{L}_\tau + \mathcal{L}_{\mathcal{L}_{\mathcal{L}_\tau\tau}\tau} + \mathcal{L}_{\mathcal{L}_\tau^2\tau})\gamma + \dots$$

Remark 3.16. *One is also interested in finding numerical methods that preserve modified measures. For example for quadratic ODEs, the Kahan-Hirota-Kimura discretization [25, 22, 23] can preserve some modified measures [14, 7]. In this context, one searches for aromatic Lagrangians $\mu \in \Omega_0$ and modified vector field represented by $\tau \in \Omega_1$ such that $(1 + \mu)\tau$ is solenoidal.*

Let us adapt Noether's theory in the context of volume-preservation. While Theorem 3.14 and Remark 3.15 focus on Lagrangians in Ω_0 and source forms in \mathcal{I}_1 in variational calculus, we are also interested in symmetries and conservation laws on Ω_1 and Ω_2 . The following result, derived from Theorem 2.5, gives necessary conditions on the form of the modified vector field of a volume-preserving method. We recall that as the aromatic bicomplex is exact, any solenoidal form $\gamma \in \Omega_1$ is the image of a form in $\hat{\gamma} \in \Omega_2$, that is, $\gamma = d_H\hat{\gamma}$.

Theorem 3.17. *Let $\gamma = d_H\hat{\gamma} \in \Omega_1$ be solenoidal, then any aromatic vector field $\tau \in \Omega_1$ is a divergence symmetry for γ . Moreover, τ is a divergence symmetry of $\hat{\gamma}$ (and thus a symmetry of γ) if and only if there exists $\eta \in \Omega_3$ such that $H_\tau\gamma = d_H\eta$.*

There is no known non-trivial modified vector field of a volume-preserving method that has symmetries to the best of our knowledge. The better understanding of the Lie derivative, the symmetries, and especially the solenoidal vector fields would give insight on the form of a volume-preserving method and help describe the degrees of freedom we have in the choice of the method. There exists a vast literature on the description of symmetries in variational calculus, which further motivates collaborations on the application of the tools of variational calculus for the creation of volume-preserving integrators. In addition, extending the aromatic bicomplex and the aromatic Noether's theory for the study of volume-preservation on manifolds or for the exact numerical preservation of the invariant measure of ergodic stochastic systems is matter for future work.

Acknowledgements. The work of Adrien Laurent was supported by the Research Council of Norway through project 302831 "Computational Dynamics and Stochastics on Manifolds" (CODYSMA). The author would like to thank E. Bronasco, D. Manchon, H. Munthe-Kaas, and the participants of the MAGIC workshops for the enlightening discussions we shared.

References

- [1] A. Abdulle, G. Vilmart, and K. C. Zygalakis. Long time accuracy of Lie-Trotter splitting methods for Langevin dynamics. *SIAM J. Numer. Anal.*, 53(1):1–16, 2015.
- [2] I. M. Anderson. The variational bicomplex. Unpublished, 1989.
- [3] I. M. Anderson. Introduction to the variational bicomplex. In *Mathematical aspects of classical field theory (Seattle, WA, 1991)*, volume 132 of *Contemp. Math.*, pages 51–73. Amer. Math. Soc., Providence, RI, 1992.
- [4] V. I. Arnold. *Mathematical methods of classical mechanics*, volume 60 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1989. Translated from the Russian by K. Vogtmann and A. Weinstein.
- [5] V. I. Arnold, V. V. Kozlov, and A. I. Neishtadt. *Mathematical aspects of classical and celestial mechanics*, volume 3 of *Encyclopaedia of Mathematical Sciences*. Springer-Verlag, Berlin, third edition, 2006. [Dynamical systems. III], Translated from the Russian original by E. Khukhro.
- [6] G. Bogfjellmo. Algebraic structure of aromatic B-series. *J. Comput. Dyn.*, 6(2):199–222, 2019.
- [7] G. Bogfjellmo, E. Celledoni, R. McLachlan, B. Owren, and R. Quispel. Using aromas to search for preserved measures and integrals in Kahan’s method. *Submitted*, 2022.
- [8] E. Bronasco. Exotic B-series and S-series: algebraic structures and order conditions for invariant measure sampling. *Submitted*, 2022.
- [9] E. Bronasco and A. Laurent. The Hopf algebra structures of composition and substitution of exotic clumped S-series. *In preparation*, 2023.
- [10] J. C. Butcher. An algebraic theory of integration methods. *Math. Comp.*, 26:79–106, 1972.
- [11] J. C. Butcher. *Numerical methods for ordinary differential equations*. John Wiley & Sons, Ltd., Chichester, third edition, 2016.
- [12] J. C. Butcher. *B-series: algebraic analysis of numerical methods*. Springer, 2021.
- [13] D. Calaque, K. Ebrahimi-Fard, and D. Manchon. Two interacting Hopf algebras of trees: a Hopf-algebraic approach to composition and substitution of B-series. *Adv. in Appl. Math.*, 47(2):282–308, 2011.
- [14] E. Celledoni, C. Evripidou, D. I. McLaren, B. Owren, G. R. W. Quispel, and B. K. Tapley. Detecting and determining preserved measures and integrals of birational maps. *Journal of Computational Dynamics*, 2022.
- [15] P. Chartier, E. Hairer, and G. Vilmart. Algebraic structures of B-series. *Found. Comput. Math.*, 10(4):407–427, 2010.
- [16] P. Chartier and A. Murua. Preserving first integrals and volume forms of additively split systems. *IMA J. Numer. Anal.*, 27(2):381–405, 2007.

- [17] A. Connes and D. Kreimer. Hopf algebras, renormalization and noncommutative geometry. *Comm. Math. Phys.*, 199(1):203–242, 1998.
- [18] G. Fløystad, D. Manchon, and H. Z. Munthe-Kaas. The universal pre-Lie-Rinehart algebras of aromatic trees. In *Geometric and harmonic analysis on homogeneous spaces and applications*, volume 366 of *Springer Proc. Math. Stat.*, pages 137–159. Springer, Cham, [2021] ©2021.
- [19] E. Hairer. Backward error analysis for multistep methods. *Numer. Math.*, 84(2):199–232, 1999.
- [20] E. Hairer, C. Lubich, and G. Wanner. *Geometric numerical integration*, volume 31 of *Springer Series in Computational Mathematics*. Springer-Verlag, Berlin, second edition, 2006. Structure-preserving algorithms for ordinary differential equations.
- [21] E. Hairer and G. Wanner. On the Butcher group and general multi-value methods. *Computing (Arch. Elektron. Rechnen)*, 13(1):1–15, 1974.
- [22] R. Hirota and K. Kimura. Discretization of the Euler top. *J. Phys. Soc. Japan*, 69(3):627–630, 2000.
- [23] R. Hirota and K. Kimura. Discretization of the Lagrange top. *J. Phys. Soc. Japan*, 69(10):3193–3199, 2000.
- [24] A. Iserles, G. R. W. Quispel, and P. S. P. Tse. B-series methods cannot be volume-preserving. *BIT Numer. Math.*, 47(2):351–378, 2007.
- [25] W. Kahan. Unconventional numerical methods for trajectory calculations. *Unpublished lecture notes*, 1:13, 1993.
- [26] F. Kang and Z. J. Shang. Volume-preserving algorithms for source-free dynamical systems. *Numer. Math.*, 71(4):451–463, 1995.
- [27] A. Laurent. *Algebraic Tools and Multiscale Methods for the Numerical Integration of Stochastic Evolutionary Problems*. PhD thesis, University of Geneva, 2021.
- [28] A. Laurent, R. I. McLachlan, H. Z. Munthe-Kaas, and O. Verdier. The aromatic bicomplex for the description of divergence-free aromatic forms and volume-preserving integrators. *To appear in Forum of Mathematics Sigma*, arXiv:2301.10998, 2023.
- [29] A. Laurent and H. Z. Munthe-Kaas. The universal equivariance properties of exotic aromatic B-series. *Submitted*, arXiv:2305.10993, 2023.
- [30] A. Laurent and G. Vilmart. Exotic aromatic B-series for the study of long time integrators for a class of ergodic SDEs. *Math. Comp.*, 89(321):169–202, 2020.
- [31] A. Laurent and G. Vilmart. Order conditions for sampling the invariant measure of ergodic stochastic differential equations on manifolds. *Found. Comput. Math.*, 22(3):649–695, 2022.
- [32] J. M. Lee. *Introduction to smooth manifolds*, volume 218 of *Graduate Texts in Mathematics*. Springer, New York, second edition, 2013.
- [33] A. Lejay. Constructing general rough differential equations through flow approximations. *Electron. J. Probab.*, 27:Paper No. 7, 24, 2022.

- [34] D. Manchon and A. Saïdi. Lois pré-Lie en interaction. *Comm. Algebra*, 39(10):3662–3680, 2011.
- [35] R. I. McLachlan, K. Modin, H. Munthe-Kaas, and O. Verdier. B-series methods are exactly the affine equivariant methods. *Numer. Math.*, 133(3):599–622, 2016.
- [36] R. I. McLachlan, K. Modin, H. Munthe-Kaas, and O. Verdier. Butcher series: a story of rooted trees and numerical methods for evolution equations. *Asia Pac. Math. Newsl.*, 7(1):1–11, 2017.
- [37] H. Munthe-Kaas and O. Verdier. Aromatic Butcher series. *Found. Comput. Math.*, 16(1):183–215, 2016.
- [38] H. Z. Munthe-Kaas and K. K. Føllesdal. Lie-Butcher series, geometry, algebra and computation. In *Discrete mechanics, geometric integration and Lie-Butcher series*, volume 267 of *Springer Proc. Math. Stat.*, pages 71–113. Springer, Cham, 2018.
- [39] E. Noether. Invariant variation problems. *Transport Theory Statist. Phys.*, 1(3):186–207, 1971.
- [40] P. J. Olver. *Applications of Lie groups to differential equations*, volume 107 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1993.
- [41] J.-M. Oudom and D. Guin. On the Lie enveloping algebra of a pre-Lie algebra. *J. K-Theory*, 2(1):147–167, 2008.
- [42] L. Rahm. An operadic approach to substitution in Lie-Butcher series. *Forum Math. Sigma*, 10:Paper No. e20, 29, 2022.
- [43] A. Saïdi. On a pre-Lie algebra defined by insertion of rooted trees. *Lett. Math. Phys.*, 92(2):181–196, 2010.
- [44] A. Saïdi. *Algèbres de Hopf d’arbres et structures pré-Lie*. PhD thesis, Université Blaise Pascal-Clermont-Ferrand II, 2011.